

Reference Numbers are from “Discrimination of Chiral Compounds Using NMR Spectroscopy,” by Thomas J. Wenzel, Wiley Press, 2007

Aryl Methoxy Acetic Acids (General review articles)

321. Seco JM, Quiñoá E, Riguera R. A practical guide for the assignment of the absolute configuration of alcohols, amines and carboxylic acids by NMR. *Tetrahedron: Asymmetry* **2001**, 12, 2915-2925.
322. Seco JM, Quiñoá E, Riguera R. The assignment of absolute configuration by NMR. *Chem. Rev.* **2004**, 104, 17-117.

Modified Mosher Method

17. Kusumi T, Fujita Y, Ohtani I, Kakisawa H. Anomaly in the modified Mosher's method: absolute configurations of some marine cembranolides. *Tetrahedron Lett.* **1991**, 32, 2923-2926.
18. Ohtani I, Kusumi T, Kashman Y, Kakisawa H. High-field FT NMR application of Mosher's method. The absolute configuration of marine terpenoids. *J. Am. Chem. Soc.* **1991**, 113, 4092-4096.
19. Kusumi T, Ohtani, II. Determination of the absolute configuration of biologically active compounds by the modified Mosher's method. *Biology-Chemistry Interface*, **1999**, 103-137.

Synthesis of MPA derivatives

149. Trost BM, Belletire JL, Godleski S, McDougal PG, Balkovec JM. On the use of the O-methylmandelate ester for establishment of absolute configuration of secondary alcohols. *J. Org. Chem.* **1986**, 51, 2370-2374.
139. Porto S, Durán J, Seco JM, Quiñoá E, Riguera R. “Mix and shake” method for configurational assignment by NMR: application to chiral amines and alcohols. *Org. Lett.* **2003**, 5, 2979-2982.

MPA – Variable temperature method

182. Seco JM, Latypov S, Quiñoá E, Riguera R. Determination of the absolute configuration of alcohols by low temperature ¹H NMR of aryl(methoxy)acetates. *Tetrahedron: Asymmetry* **1995**, 6, 107-110.
184. Latypov SK, Seco JM, Quiñoá E, Riguera R. Are both the (R)- and the (S)-MPA esters really needed for the assignment of the absolute configuration of secondary alcohols by NMR? The use of a single derivative. *J. Am. Chem. Soc.* **1998**, 120, 877-882.

MPA – Barium method

134. García R, Seco JM, Vázquez SA, Quiñoá E, Riguera R. Absolute configuration of secondary alcohols by ¹H NMR: in situ complexation of α -methoxyphenylacetic acid esters with barium(II). *J. Org. Chem.* **2002**, 67, 4579-4589.
187. López B, Quiñoá E, Riguera R. Complexation with barium(II) allows the inference of the absolute configuration of primary amines by NMR. *J. Am. Chem. Soc.* **1999**, 121, 9724-9725.
188. Garcia R, Seco JM, Vázquez SA, Quiñoá E, Riguera R. Role of barium(II) in the determination of the absolute configuration of chiral amines by ¹H NMR spectroscopy. *J. Org. Chem.* **2006**, 71, 1119-1130.

MTPA – Addition of La(hfa)₃

135. Omata K, Fujiwara T, Kabuto K. Use of diamagnetic lanthanide complex for extending the scope of NMR determination of absolute configuration by the modified Mosher's method. *Tetrahedron: Asymmetry* **2002**, *13*, 1655-1662.

9-AMA/2-AMA

14. Latypov SK, Seco JM, Quiñoá E, Riguera R. Conformational structure and dynamics of arylmethoxyacetates: DNMR spectroscopy and aromatic shielding effect. *J. Org. Chem.* **1995**, *60*, 504-515.

15. Seco JM, Latypov SK, Quiñoá E, Riguera R. Determining factors in the assignment of the absolute configuration of alcohols by NMR. The use of anisotropic effects on remote positions. *Tetrahedron* **1997**, *53*, 8541-8564.

307. Kouda K, Kusumi T, Ping X, Kan Y, Hashimoto T, Asakawa Y. 2-anthrylmethoxyacetic acid, a new chiral anisotropic reagent for elucidating the absolute configuration of acyclic alcohols. *Tetrahedron Lett.* **1996**, *37*, 4541-4544.

308. Seco JM, Quiñoá E, Riguera R. 9-Anthrylmethoxyacetic acid esterification shifts-correlation with the absolute stereochemistry of secondary alcohols. *Tetrahedron* **1999**, *55*, 569-584.

309. Kouda K, Ooi T, Kaya K, Kusumi T. Absolute stereostructure of a 2,3,7,13-tetrahydroxyoctadecanoic acid, the framework of taurolipid B produced by a fresh-water protozoan, *Tetrahymena thermophila*. *Tetrahedron Lett.* **1996**, *37*, 6347-6350.

MαNP

287. Kasai Y, Naito J, Kuwahara S, Watanabe M, Ichikawa A, Harada N. Novel chiral molecular tools for preparation of enantiopure alcohols by resolution and simultaneous determination of their absolute configurations by the ¹H NMR anisotropy method. *J. Synth. Org. Chem., Jpn.* **2004**, *62*, 48-61.

288. Fujita T, Kuwahara S, Watanabe M, Harada N. Crystalline state conformation of 2-methoxy-2-(1-naphthyl)propionic acid ester. *Enantiomer* **2002**, *7*, 219-223.

289. Taji H, Watanabe M, Harada N, Naoki H, Ueda Y. Diastereomer method for determining % ee by ¹H NMR and/or MS spectrometry with complete removal of the kinetic resolution effect. *Org. Lett.* **2002**, *4*, 2699-2702.

290. Ichikawa A. Application of αMNPA to stereochemical studies of monoterpene alcohol. *Chirality* **1999**, *11*, 70-74.

291. Taji H, Kasai Y, Sugio A, Kuwahara S, Watanabe M, Harada N, Ichikawa A. Practical enantioresolution of alcohols with 2-methoxy-2-(1-naphthyl)propionic acid and determination of their absolute configurations by the ¹H NMR anisotropy method. *Chirality* **2002**, *14*, 81-84.

292. Kasai Y, Watanabe M, Harada N. Convenient method for determining the absolute configuration of chiral alcohols with racemic ¹H NMR anisotropy reagent, MαNP acid: use of HPLC-CD detector. *Chirality* **2003**, *15*, 295-299.

293. Kosaka M, Sekiguchi S, Naito J, Uemura M, Kuwahara S, Watanabe M, Harada N, Hiroi K. Synthesis of enantiopure phthalides including 3-butylphthalide, a fragrance component of celery oil, and determination of their absolute configurations. *Chirality* **2005**, *17*, 218-232.

294. Kosaka M, Sugito T, Kasai Y, Kuwahara S, Watanabe M, Harada N, Job GE, Shvet A, Pirkle WH. Enantioresolution and absolute configurations of chiral meta-substituted diphenylmethanols as determined by X-ray crystallographic and ¹H NMR anisotropy methods. *Chirality* **2003**, *15*, 324-328.

295. Kasai Y, Taji H, Fujita T, Yamamoto Y, Akagi M, Sugio A, Kuwahara S, Watanabe M, Harada N, Ichikawa A, Schurig V. M α NP acid, a powerful chiral molecular tool for preparation of enantiopure alcohols by resolution and determination of their absolute configurations by the ^1H NMR anisotropy method. *Chirality* **2004**, 16, 569-585.

296. Harada N, Watanabe M, Kuwahara S, Sugio A, Kasai Y, Ichikawa A. 2-Methoxy-2-(1-naphthyl)propionic acid, a powerful chiral auxiliary for enantioresolution of alcohols and determination of their absolute configurations by the ^1H NMR anisotropy method. *Tetrahedron: Asymmetry* **2000**, 11, 1249-1253.

CFTA

249. Takeuchi Y, Konishi M, Hori H, Takahashi T, Kometani T, Kirk KL. Efficient synthesis of a new, highly versatile chiral derivatizing agent, α -cyano- α -fluoro-*p*-tolylacetic acid (CFTA). *Chem. Commun.* **1998**, 365-366.

250. Takahashi T, Fukuishima A, Tanaka Y, Takeuchi Y, Kabuto K, Kabuto C. CFTA, a new efficient agent for determination of absolute configurations of chiral secondary alcohols. *Chem. Commun.* **2000**, 788-789.

251. Fujiwara T, Omata K, Kabuto K, Kabuto C, Takahashi T, Segawa M, Takeuchi Y. Determination of the absolute configurations of α -amino esters from the ^{19}F NMR chemical shifts of their CFTA amide diastereomers. *Chem. Commun.* **2001**, 2694-2695.

252. Fujiwara T, Sasaki M, Omata K, Kabuto C, Kabuto K, Takeuchi Y. An efficient procedure for the resolution of α -cyano- α -fluoro-*p*-tolylacetic acid (CFTA) via the diastereomeric N-carbobenzyloxy-*cis*-1-amino-2-indanol esters. *Tetrahedron: Asymmetry* **2004**, 15, 555-563.

253. Takeuchi Y, Fujisawa H, Noyori R. A very reliable method for determination of absolute configuration of chiral secondary alcohols by ^1H NMR spectroscopy. *Org. Lett.* **2004**, 6, 4607-4610.

MPA (Diols)

171. Seco JM, Martino M, Quiñoá E, Riguera R. Absolute configuration of 1, n-diols by NMR: the importance of the combined anisotropic effects in bis-arylmethoxyacetates. *Org. Lett.* **2000**, 2, 3261-3264.

172. Freire F, Seco JM, Quiñoá E, Riguera R. Determining the absolute stereochemistry of secondary/secondary diols by ^1H NMR: basis and applications. *J. Org. Chem.* **2005**, 70, 3778-3790.

MTPA (Primary alcohols)

82. Tsuda M, Toriyabe Y, Endo T, Kobayashi J. Application of modified Mosher's method for primary alcohols with a methyl group at C2 position. *Chem. Pharm. Bull.* **2003**, 51, 448-451.

9-AMA (Primary alcohols)

313. Ferreiro MJ, Latypov SK, Quiñoá E, Riguera R. Determination of the absolute configuration and enantiomeric purity of chiral primary alcohols by ^1H NMR of 9-anthrylmethoxyacetates. *Tetrahedron: Asymmetry* **1996**, 7, 2195-2198.

314. Latypov SK, Ferreiro MJ, Quiñoá E, Riguera R. Assignment of the absolute configuration of β -chiral primary alcohols by NMR: scope and limitations. *J. Am. Chem. Soc.* **1998**, 120, 4741-4751.

MTPA (Tertiary alcohols)

87. Izumi S, Moriyoshi H, Hirata T. Identification of absolute configuration of tertiary alcohols by combination of Mosher's method and conformational analysis. *Bull. Chem. Soc. Jpn.* **1994**, 67, 2600-2602.

276. Takahashi H, Kato N, Iwashima M, Iguchi K. Determination of absolute configurations of tertiary alcohols by NMR spectroscopy. *Chem. Lett.* **1999**, 1181-1182.

MTPA (Secondary amines)

88. Hoyer TR, Renner MK. MTPA (Mosher) amides of cyclic secondary amines: conformational aspects and a useful method for assignment of amine configuration. *J. Org. Chem.* **1996**, 61, 2056-2064.

89. Hoyer TR, Renner MK. Applications of MTPA (Mosher) amides of secondary amines: assignment of absolute configuration in chiral cyclic amines. *J. Org. Chem.* **1996**, 61, 8489-8495.

(MTPA (Primary amines)

95. Seco JM, Latypov SK, Quiñoá E, Riguera R. Choosing the right reagent for the determination of the absolute configuration of amines by NMR: MTPA or MPA? *J. Org. Chem.* **1997**, 62, 7569-7574.

96. Kusumi T, Fukushima T, Ohtani I, Kakisawa H. Elucidation of the absolute configurations of amino acids and amines by the modified Mosher's method. *Tetrahedron Lett.* **1991**, 32, 2939-2942.

BPG (Primary amines)

255. Seco JM, Quiñoá E, Riguera R. Boc-phenylglycine: the reagent of choice for the assignment of the absolute configuration of α -chiral primary amines by ^1H NMR spectroscopy. *J. Org. Chem.* **1999**, 64, 4669-4675.

256. Ito Y, Ishida K, Okada S, Murakami M. The absolute stereochemistry of anachelins, siderophores from the cyanobacterium *Anabaena cylindrica*. *Tetrahedron* **2004**, 60, 9075-9080.

Camphanic Acid

327. Furukawa J, Iwasaki S, Okuda S. Method for determination of enantiomeric composition and absolute configuration of 2,3-deuterated 3-alkylpropanols. *Tetrahedron Lett.* **1983**, 24, 5257-5260.

330. Fretz H, Woggon W, Voges R. 49. The allylic oxidation of geraniol catalyzed by cytochrome P-450_{Cath.}, proceeding with retention of configuration. *Helv. Chim. Acta* **1989**, 72, 391-400.

331. Ravn MM, Jin Q, Coates RM. Synthesis of allylic isoprenoid diphosphates by $\text{S}_{\text{N}}2$ displacement of diethyl phosphate. *Eur. J. Org. Chem.* **2000**, 1401-1410.

334. Armarego WLF, Milloy BA, Pendergast W. A highly stereospecific synthesis of (R)- and (S)-[2- $^2\text{H}_1$]glycine. *J. Chem. Soc., Perkin Trans. 1* **1976**, 2229-2237.

335. Pontoni G, Coward JK, Orr GR, Gould SJ. Stereochemical studies of enzyme-catalyzed alkyl-transfer reactions. An NMR method for distinguishing between the two prochiral hydrogens at C-1' of spermidine. *Tetrahedron Lett.* **1983**, 24, 151-154.

336. Parker D, Taylor RJ, Ferguson G, Tonge A. Origins of the proton NMR chemical shift non-equivalence in the diastereotopic methylene protons of camphanamides. *Tetrahedron* **1986**, 42, 617-622.

337. Williams RM, Zhai D, Sinclair PJ. Asymmetric synthesis of (R)- and (S)-[2- $^2\text{H}_1$]glycine. *J. Org. Chem.* **1986**, 51, 5021-5022.

338. Ramer SE, Cheng H, Vederas JC. Investigations of polypeptide biosynthesis: formation of peptide amides. *Pure Appl. Chem.* **1989**, 61, 489-492.

339. Hameršak Z, Selestrin A, Lesac A, Šunjić V. Conformational study of α -arylethylamides of (–)-camphanic acid. *Tetrahedron: Asymmetry* **1998**, 9, 1891-1897.
340. Brown JM, Parker D. Determination of optical purity at isotopically chiral sites by ^2H NMR. *Tetrahedron Lett.* **1981**, 22, 2815-2818.
341. Parker D. ^1H and ^2H Nuclear magnetic resonance determination of the enantiomeric purity and absolute configuration of α -deuteriated primary carboxylic acids, alcohols, and amines. *J. Chem. Soc., Perkin Trans. 2* **1983**, 83-88.

2-(2,3-Anthracenedicarboximido)cyclohexane carboxylic acid

360. Akasaka K, Imaizumi K, Sakakibara R, Terashima H, Ohru H. Chiral discrimination of branched alkyl chain by labeling with chiral derivatization reagents. *Tennen Yuki Kagobutsu Toronkai Koen Yoshishu* **2000**, 42, 565-570.
361. Ohru H, Terashima H, Imaizumi K, Akasaka K. A solution of the “intrinsic problem” of diastereomer method in chiral discrimination. Development of a method for highly efficient and sensitive discrimination of chiral alcohols. *Proc. Japan Acad. Ser. B Phys. Biol. Sci.* **2002**, 78, 69-72.
362. Imaizumi K, Terasima H, Akasaka K, Ohru H. Highly potent chiral labeling reagents for the discrimination of chiral alcohols. *Anal. Sci.* **2003**, 19, 1243-1249.
363. Ohtaki T, Akasaka K, Kabuto C, Ohru H. Chiral discrimination of secondary alcohols by both ^1H -NMR and HPLC after labeling with a chiral derivatization reagent, 2-(2,3-anthracenedicarboximide)cyclohexane carboxylic acid. *Chirality* **2005**, 17, S171-S176.

Pirkle's Alcohol (TFPE/TFAE)

408. Pirkle WH, Burlingame TG, Beare SD. Optically active NMR solvents VI. The determination of optical purity and absolute configuration of amines. *Tetrahedron Lett.* **1968**, 5849-5852.
409. Pirkle WH, Beare SD. Optically active solvents in nuclear magnetic resonance spectroscopy. VII. Direct determination of optical purities and correlation of absolute configurations of sulfoxides. *J. Am. Chem. Soc.* **1968**, 90, 6250-6251.
410. Pirkle WH, Pavlin MS. Determination of the enantiomeric purity of compound chiral by virtue of C-13 labelling. *J. Chem. Soc., Chem. Commun.* **1974**, 274-275.
411. Pirkle WH, Beare SD, Muntz RL. Assignment of absolute configuration of sulfoxides by NMR. A solvation model. *Tetrahedron Lett.* **1974**, 2295-2298.
415. Pirkle WH, Beare SD, Muntz RL. Optically active solvents for nuclear magnetic resonance. X. Enantiomeric nonequivalence of sulfinamides, sulfinates, sulfites, thiosulfinates, phosphine oxides, and amine oxides. *J. Am. Chem. Soc.* **1969**, 91, 4575.
419. Pirkle WH, Muntz RL, Paul IC. Chiral nuclear magnetic resonance solvents. XI. A method for determining the absolute configuration of chiral N,N-dialkylarylamine oxides. *J. Am. Chem. Soc.* **1971**, 93, 2817-2819.
420. Pirkle WH, Beare SD. Optically active solvents in nuclear magnetic resonance spectroscopy. IX. Direct determinations of optical purities and correlations of absolute configurations of α -amino acids. *J. Am. Chem. Soc.* **1969**, 91, 5150-5155.
422. Forni A, Moretti I, Torre G. Absolute configuration at chiral nitrogen in oxaziridines configurational correlations by NMR spectroscopy in optically active solvating agents. *Tetrahedron Lett.* **1978**, 2941-2944.

423. Bucciarelli M, Forni A, Moretti I, Torre G. Asymmetric synthesis at nitrogen by oxidation of imines with m-chloroperbenzoic acid in the presence of optically active carbinols. Absolute stereochemistry of chiral alcohol-imine-peracid solvates. *J. Chem. Soc., Perkin Trans. 1* **1980**, 2152-2161.
432. Pirkle WH, Sikkenga DL, Pavlin MS. Nuclear magnetic resonance determination of enantiomeric composition and absolute configuration of γ -lactones using chiral 2,2,2-trifluoro-1-(9-anthryl)ethanol. *J. Org. Chem.* **1977**, *42*, 384-387.
433. Pirkle WH, Sikkenga DL. The use of chiral solvating agents for nuclear magnetic resonance determination of enantiomeric purity and absolute configuration of lactones. Consequences of three-point interactions. *J. Org. Chem.* **1977**, *42*, 1370-1374.
437. Pirkle WH, Robertson MR, Hyun MH. A liquid chromatographic method for resolving chiral lactams as their diastereomeric ureide derivatives. *J. Org. Chem.* **1984**, *49*, 2433-2437.

2-(2,3-Anthracenedicarboximido)-1-cyclohexanol

501. Akasaka K, Imaizumi K, Shichijyukari S, Ohru H, Kabuto C. Enantiomeric discrimination of carboxylic acids having chiral centers remote from the carboxyl group by labeling with chiral fluorescent derivatization reagents. *Tennen Yuki Kagobutsu Toronkai Koen Yoshishu* **1998**, *40*, 79-84.

Methyl mandelate

332. Brown JM, Parker D. Mechanism of asymmetric homogeneous hydrogenation. Rhodium-catalyzed reductions with deuterium and hydrogen deuteride. *Organometallics* **1982**, *1*, 950-956.
341. Parker D. ^1H and ^2H Nuclear magnetic resonance determination of the enantiomeric purity and absolute configuration of α -deuteriated primary carboxylic acids, alcohols, and amines. *J. Chem. Soc., Perkin Trans. 2* **1983**, 83-88.
511. Schwab JM, Klassen JB. Steric course of the allylic rearrangement catalyzed by β -hydroxydecanoylthioester dehydrase. Mechanistic implications. *J. Am. Chem. Soc.* **1984**, *106*, 7217-7227.

Menthol

526. Lewis RA, Korpiun O, Mislow K. Configurational correlation of phosphinates by nuclear magnetic resonance and optical rotatory dispersion. *J. Am. Chem. Soc.* **1967**, *89*, 4786-4787.
527. Lewis RA, Korpiun O, Mislow K. Proton magnetic resonance spectra of menthyl phosphinates. *J. Am. Chem. Soc.* **1968**, *90*, 4847-4853.
528. Vitharana D, France JE, Scarpetti D, Bonneville GW, Majer P, Tsukamoto T. Synthesis and biological evaluation of (R)- and (S)-2-(phosphonomethyl)pentanedioic acids as inhibitors of glutamate carboxypeptides II. *Tetrahedron: Asymmetry* **2002**, *13*, 1609-1614.
529. Holt A, Jarvie AWP, Jarvis GJ. The preparation and determination of optical purity of methylphenyl-1-naphthyl(-)-menthoxyasilanes. *J. Organomet. Chem.* **1970**, *21*, 75-77.
530. Holt A, Jarvie AWP, Jarvis GJ. Preparation and spectroscopic properties of some new asymmetric organosilanes. *J. Chem. Soc., Perkin Trans. 2* **1973**, 114-122.
531. Corriu RJP, Lanneau GF. Synthèse et structure de nouveaux organosilanes chiraux bifonctionnels. *J. Organomet. Chem.* **1974**, *64*, 63-78.

536. Ichikawa A, Hiradate S, Sugio A, Kuwahara S, Watanabe M, Harada N. Absolute configuration of 2-hydroxy-2-(1-naphthyl)propionic acid as determined by the ^1H NMR anisotropy method. *Tetrahedron: Asymmetry* **1999**, *10*, 4075-4078.

537. Ichikawa A, Hiradate S, Sugio A, Kuwahara S, Watanabe M, Harada N. Absolute configuration of 2-methoxy-2-(2-naphthyl)propionic acid as determined by the ^1H NMR anisotropy method. *Tetrahedron: Asymmetry* **2000**, *11*, 2669-2675.

Glycosidation shifts

550. Kasai R, Suzuo M, Asakawa J, Tanaka O. Carbon-13 chemical shifts of isoprenoid- β -D-glucopyranosides and - β -D-mannopyranosides. Stereochemical influences of aglycone alcohols. *Tetrahedron Lett.* **1977**, 175-178.

551. Tori K, Seo S, Yoshimura Y, Arita H, Tomita Y. Glycosidation shifts in carbon-13 NMR spectroscopy: carbon-13 signal shifts from aglycone and glucose to glucoside. *Tetrahedron Lett.* **1977**, 179-182.

552. Seo S, Tomita Y, Tori K, Yoshimura Y. Determination of the absolute configuration of a secondary hydroxy group in a chiral secondary alcohol using glycosidation shifts in carbon-13 nuclear magnetic resonance spectroscopy. *J. Am. Chem. Soc.* **1978**, *100*, 3331-3339.

553. Horibe I, Seo S, Yoshimura Y, Tori K. Glucosidation shifts of allylic and benzylic alcohols in ^{13}C NMR spectroscopy. *Org. Magn. Reson.* **1984**, *22*, 428-430.

554. Faghih R, Fontaine C, Horibe I, Imamura PM, Lukacs G, Olesker A, Seo S. Determination of the absolute configuration of chiral secondary alcohols from tetra-O-acetylglucosidation-induced ^1H nuclear magnetic resonance shifts. *J. Org. Chem.* **1985**, *50*, 4918-4920.

555. Seroka P, Płosiński M, Czub J, Sowiński P, Pawlak J. Monosaccharides as internal probes for the determination of the absolute configuration of 2-butanol. *Magn. Reson. Chem.* **2006**, *44*, 132-138.

556. Kurokawa-Nose Y, Shimada T, Wada H, Tanaka N, Murakami T, Saiki Y. Chemical and chemotaxonomical studies of ferns. LXXXIV. A novel 2-tetralol-type xyloside from *Asplenium wilfordii*. *Chem. Pharm. Bull.* **1993**, *41*, 930-932.

565. Trujillo M, Morales EQ, Vázquez JT. Tetra-O-benzoylglucosylation: a new ^1H nuclear magnetic resonance method for determination of the absolute configuration of secondary alcohols. *J. Org. Chem.* **1994**, *59*, 6637-6642.

566. Morales EQ, Padrón JI, Trujillo M, Vázquez JT. CD and ^1H NMR study of the rotational population dependence of the hydroxymethyl group in β glucopyranosides on the aglycon and its absolute configuration. *J. Org. Chem.* **1995**, *60*, 2537-2548.

Fuco/arabinofuranoside

575. Kobayashi M. The fucofuranoside method, a new ^1H and ^{13}C nuclear magnetic resonance method to determine the absolute configuration of secondary alcohols. *Tetrahedron* **1997**, *53*, 5973-5994.

576. Ma Z, Hano Y, Qiu F, Chen Y, Nomura T. Determination of the absolute stereochemistry of lupine triterpenoids by fucofuranoside method and ORD spectrum. *Tetrahedron Lett.* **2004**, *45*, 3261-3263.

577. Kobayashi M. The fucofuranoside method for determining the absolute configuration of the tertiary alcohols substituted with methyl and two methylene groups. *Tetrahedron* **1998**, *54*, 10987-10998.

578. Kobayashi M. The ^{13}C NMR method for determining the absolute configuration of the 1,2-glycols consisting of secondary and tertiary hydroxyl groups. *Tetrahedron* **2000**, *56*, 1661-1665.

579. Kobayashi M. The arabinofuranoside method, a convenient substitute of the fucofuranoside method for determining the absolute configuration of the secondary alcohols. *Tetrahedron* **2002**, 58, 9365-9371.

BINOL

582. Toda F, Mori K, Okada J, Node M, Itoh A, Oomine K, Fuji K. New chiral shift reagents, optically active 2,2'-dihydroxy-1,1'-binaphthyl and 1,6-di(o-chlorophenyl)-1,6-diphenylhexa-2,4-diyne-1,6-diol. *Chem. Lett.* **1988**, 131-134.

583. Toda F, Mori K, Satō A. A determination method of absolute configuration. *Bull. Chem. Soc. Jpn.* **1988**, 61, 4167-4169.

584. Drabowicz J, Duddeck H. ¹H NMR spectral nonequivalence of sulphoxide enantiomers in the presence of 2,2'-dihydroxy-1,1'-binaphthyl. *Sulfur Lett.* **1989**, 10, 37-40.

585. Reynolds DP, Hollerton JC, Richards SA. Optical purity determination by ¹H NMR. *Analytical Applications of Spectroscopy*. Creaser CS, Davies AMC, Ed. Royal Society of Chemistry: London, England, **1988**, 346-348.

Butane-2,3-diol

596. Helder R, Arends R, Bolt W, Hiemstra H, Wynberg H. Alkaloid catalyzed asymmetric synthesis. III. The addition of mercaptans to 2-cyclohexene-1-one; determination of enantiomeric excess using carbon-13 NMR. *Tetrahedron Lett.* **1977**, 2181-2182.

597. Hiemstra H, Wynberg H. Determination of the enantiomeric purity of chiral ketones using ¹³C NMR spectroscopy of their diastereomeric cyclic ketals. *Tetrahedron Lett.* **1977**, 2183-2186.

598. ten Hoeve W, Wynberg H. Chiral spiranes. Optical activity and nuclear magnetic resonance spectroscopy as a proof for stable twist conformations. *J. Org. Chem.* **1979**, 44, 1508-1514.

599. Meyers AI, Williams DR, Erickson GW, White S, Druelinger M. Enantioselective alkylation of ketones via chiral, nonracemic lithioenamines. An asymmetric synthesis of α -alkyl and α,α' -dialkyl cyclic ketones. *J. Am. Chem. Soc.* **1981**, 103, 3081-3087.

600. 2410.

Lemière GL, Dommissie RA, Lepoivre JA, Alderweireldt FC, Hiemstra H, Wynberg H, Jones JB, Toone EJ. Determination of the absolute configuration of six-membered-ring ketones by ¹³C NMR. *J. Am. Chem. Soc.* **1987**, 109, 1363-1370.

PEA/NEA/AEA

611. Pirkle WH, Beare SD. Nonequivalence of the nuclear magnetic resonance spectra of enantiomers in optically active solvents. IV. Assignment of absolute configuration. *J. Am. Chem. Soc.* **1967**, 89, 5485-5487.

623. Helmchen G, Ott R, Sauber K. Objective separation and absolute configuration of enantiomeric carboxylic acids and amines. 1. *Tetrahedron Lett.* **1972**, 3873-3878.

624. Hoyer TR, Koltun DO. An NMR strategy for determination of configuration of remote stereogenic centers: 3-methylcarboxylic acids. *J. Am. Chem. Soc.* **1998**, 120, 4638-4643.

625. Hoyer TR, Hamad AS, Koltun DO, Tennakoon MA. An NMR method for determination of configuration of β -substituted carboxylic acids. *Tetrahedron Lett.* **2000**, 41, 2289-2293.

631. Ghislandi V, La Manna A, Azzolina O, Gazzaniga A, Vercesi D. Configurational relationships in antiphlogistic hydratropic acids. *Farmaco* **1982**, 37, 81-93.

635. Mikołajczyk M, Para M, Ejchart A, Jurczak J. Correlation of the absolute configuration of phosphorus thioacids by nuclear magnetic resonance spectroscopy. *Chem. Commun.* **1970**, 654.
636. Mikołajczyk M, Omelańczuk J, Leitloff M, Drabowicz J, Ejchart A, Jurczak J. ^1H , ^{31}P , and ^{13}C nuclear magnetic resonance nonequivalence of diastereomeric salts of chiral phosphorus thio acids with optically active amines. A method for determining the optical purity and configuration of chiral phosphorus thio acids. *J. Am. Chem. Soc.* **1978**, *100*, 7003-7008.
637. Głowacki Z, Hoffman M, Rachoń J. Direct ^{31}P NMR determination of the enantiomeric composition of 1-hydroxyalkylphosphonates using 1-(1-naphthyl)ethylamine as a chiral agent. *Phosphorus, Sulfur Silicon Relat. Elem.* **1993**, *82*, 39-47.
640. Hoyer G, Rosenberg D, Rufer C, Seeger A. Bestimmung der optischen reinheit und absoluten konfiguration von enantiomeren aralkylaminen. *Tetrahedron Lett.* **1972**, 985-988.
641. Takano S, Kasahara C, Ogasawara K. A simultaneous determination of enantiomeric purity and chirality of a certain cyclic enone relating to the gibbane framework. *Heterocycles* **1981**, *16*, 1669-1671.
642. Mamlok L, Marquet A, Lacombe L. Feasibility of determining the optical purity of ketones by NMR. *Tetrahedron Lett.* **1971**, 1039-1042.
644. Nowick JS, Powell NA, Nguyen TM, Noronha G. An improved method for the synthesis of enantiomerically pure amino acid ester isocyanates. *J. Org. Chem.* **1992**, *57*, 7364-7366.

PGME

36. Sata NU, Wada S, Matsunaga S, Watabe S, van Soest RWM, Fusetani N. Rubrosides A-H, new bioactive tetramic acid glycosides from the marine sponge *Siliquariaspongia japonica*. *J. Org. Chem.* **1999**, *64*, 2331-2339.
673. Nagai Y, Kusumi T. New chiral anisotropic reagents for determining the absolute configuration of carboxylic acids. *Tetrahedron Lett.* **1995**, *36*, 1853-1856.
674. Yabuuchi T, Ooi T, Kusumi T. Application of phenylglycine methyl ester (PGME) to determination of the absolute configuration of carboxylic acids having phenylalkyl group. *Chirality* **1997**, *9*, 550-555.
675. Yabuuchi T, Kusumi T. Phenylglycine methyl ester, a useful tool for absolute configuration determination of various chiral carboxylic acids. *J. Org. Chem.* **2000**, *65*, 397-404.

1,2-Diphenyl-1,2-diaminoethane

740. Alexakis A, Frutos JC, Mangeney P. An easy and fast way to determine the enantiomeric purity of substituted cyclanones. *Tetrahedron: Asymmetry* **1993**, *4*, 2431-2434.
741. Mangeney P, Alexakis A, Normant JF. Resolution and determination of enantiomeric excesses of chiral aldehydes via chiral imidazolidines. *Tetrahedron Lett.* **1988**, *29*, 2677-2680.
742. Cuvilot D, Mangeney P, Alexakis A, Normant J, Lellouche J. Chiral trifluoro diamines as convenient reagents for determining the enantiomeric purity of aldehydes by use of ^{19}F NMR spectroscopy. *J. Org. Chem.* **1989**, *54*, 2420-2425.

Amide chiral solvating agents (DNB-PEA, DNB-Leu, Whelk-O-1)

754. Deshmukh M, Duñach E, Juge S, Kagan HB. A convenient family of chiral shift reagents for measurement of enantiomeric excesses of sulfoxides. *Tetrahedron Lett.* **1984**, *25*, 3467-3470.

759. Duñach E, Kagan HB. A simple chiral shift reagent for measurement of enantiomeric excesses of phosphine oxides. *Tetrahedron Lett.* **1985**, 26, 2649-2652.
767. Pirkle WH, Tsiouras A. 3,5-Dinitrobenzoyl amino acid esters. Broadly applicable chiral solvating agents for NMR determination of enantiomeric purity. *Tetrahedron Lett.* **1985**, 26, 2989-2992.
770. Wenzel TJ, Morin CA, Brechting AA. Lanthanide-chiral resolving agent mixtures as chiral NMR shift reagents. *J. Org. Chem.* **1992**, 57, 3594-3599.
775. Pirkle WH, Gan KZ. A direct chromatographic separation of enantiomers chiral by virtue of isotopic substitution. *Tetrahedron: Asymmetry* **1997**, 8, 811-814.
776. Job GE, Shvets A, Pirkle WH, Kuwahara S, Kosaka M, Kasai Y, Taji H, Fujita K, Watanabe M, Harada N. The effects of aromatic substituents on the chromatographic enantioseparation of diarylmethyl esters with the Whelk-O1 chiral stationary phase. *J. Chromatogr. A* **2004**, 1055, 41-53.
777. Koscho ME, Pirkle WH. Investigation of a broadly applicable chiral selector used in enantioselective chromatography (Whelk-O 1) as a chiral solvating agent for NMR determination of enantiomeric composition. *Tetrahedron: Asymmetry* **2005**, 16, 3345-3351.

Phosphinothioic acid

873. Harger MJP. Nuclear magnetic resonance non-equivalence of the enantiomers in optically active samples of phosphinic amides. *J. Chem. Soc., Chem. Commun.* **1976**, 555-556.
874. Harger MJP. Proton magnetic resonance non-equivalence of the enantiomers of alkylphenylphosphinic amides. *J. Chem. Soc., Perkin Trans. 2* **1977**, 1882- 1887.
875. Ouryupin AB, Kadyko MI, Petrovskii PV, Fedin EI. Chiral discrimination in nonracemic mixtures of methanephosphonic acid, N,N'-bis(1-phenylethyl)diamides. *Chirality* **1994**, 6, 1-4.
876. Uryupin BA, Kadyko MI, Petrovskii PV, Fedin EI, Sanshe M, Wolf R, Mastryukova TA, Kabachnik MI. NMR determination of enantiomeric composition in the case of statistically controlled associate-diastereomer anisochrony. *Zh. Obshch. Khim.* **1995**, 65, 18-22.
877. Harger MJP. Chemical shift non-equivalence of enantiomers in the proton magnetic resonance spectra of partly resolved phosphinothioic acids. *J. Chem. Soc., Perkin Trans. 2* **1978**, 326-331.

P(V) reagents

895. Johnson CR, Elliott RC, Penning TD. Determination of enantiomeric purities of alcohols and amines by a ³¹P NMR technique. *J. Am. Chem. Soc.* **1984**, 106, 5020-5022.
896. Cullis PM, Iagrossi A, Rous AJ. Thiophosphoryl transfer reactions: a general synthesis and configurational analysis of O-substituted [¹⁶O, ¹⁸O]thiophosphates. *J. Am. Chem. Soc.* **1986**, 108, 7869-7870.
899. Anderson RC, Shapiro MJ. 2-Chloro-4(R),5(R)-dimethyl-2-oxo-1,3,2-dioxaphospholane, a new chiral derivatizing agent. *J. Org. Chem.* **1984**, 49, 1304-1305.
900. Brown HC, Imai, T, Desai MC, Singaram B. Chiral synthesis via organoboranes 3. Conversion of boronic esters of essentially 100% optical purity to aldehydes, acids, and homologated alcohols of very high enantiomeric purities. *J. Am. Chem. Soc.* **1985**, 107, 4980-4983.

901. Brown HC, Naik RG, Bakshi RK, Pyun C, Singaram B. Chiral synthesis via organoboranes. 4. Synthetic utility of boronic esters of essentially 100% optical purity. Synthesis of homologated boronic acids and esters of very high enantiomeric purities. *J. Org. Chem.* **1985**, 50, 5586-5592.
902. Brown HC, Kim K-W, Cole TE, Singaram B. Chiral synthesis via organoboranes. 8. Synthetic utility of boronic esters of essentially 100% optical purity. Synthesis of primary amines of very high enantiomeric purities. *J. Am. Chem. Soc.* **1986**, 108, 6761-6764.
903. Hägele G, Boisière B, Helie I, Rabiller C, Martin GJ, Martin ML. Monitoring site-specific natural hydrogen isotope fractionation using a novel phosphorus-containing chiral discriminating reagent. *Magn. Reson. Chem.* **1991**, 29, 813-822.
904. van Hoeve W, Wynberg H. The design of resolving agents. Chiral cyclic phosphoric acids. *J. Org. Chem.* **1985**, 50, 4508-4514.
905. Hulst R, Zijlstra RWJ, Feringa BL, de Vries NK, ten Hoeve W, Wynberg H. A new ^{31}P NMR method for the enantiomeric excess determination of alcohols, amines, and amino acid esters. *Tetrahedron Lett.* **1993**, 34, 1339-1342.
909. Oshikawa T, Yamashita M, Kumagai S, Seo K, Kobayashi J. Optically active diazacyclophosphamides: novel efficient reagents for the determination of the absolute configuration of amines and alcohols. *J. Chem. Soc., Chem. Commun.* **1995**, 435-436.

P(III) reagents –diamine ligands

925. Alexakis A, Mutti S, Normant JF, Mangeney P. A new reagent for a very simple and efficient determination of enantiomeric purity of alcohols by ^{31}P NMR. *Tetrahedron: Asymmetry* **1990**, 1, 437-440.
926. Alexakis A, Mutti S, Mangeney P. A new reagent for the determination of the optical purity of primary, secondary, and tertiary chiral alcohols and of thiols. *J. Org. Chem.* **1992**, 57, 1224-1237.
927. Alexakis A, Frutos JC, Mutti S, Mangeney P. Chiral diamines for a new protocol to determine the enantiomeric composition of alcohols, thiols, and amines by ^{31}P , ^1H , ^{13}C , and ^{19}F NMR. *J. Org. Chem.* **1994**, 59, 3326-3334.
928. Nowacki J, Szychowski J, Wojtasiewicz K, Matuszewska I, Jaremko M, Czarnocki Z. The determination of enantiomeric excess of the products of Baker's yeast-mediated reduction of δ -alkoxycarbonyl- β -keto esters, via the derivation to Alexakis' alkoxydiazaphosphoridine. *Heterocycles* **2002**, 57, 705-714.
929. Alexakis A, Frutos JC, Mangeney P, Meyers AI, Moorlag H. Determination of enantiomeric purity of hydroxy biaryls using ^1H and ^{31}P -NMR on their diazaphospholidine derivatives. *Tetrahedron Lett.* **1994**, 35, 5125-5128.
930. Alexakis A, Chauvin A. Determination of the enantiomeric excess of chiral carboxylic acids by ^{31}P NMR with C_2 symmetrical diamines. *Tetrahedron: Asymmetry* **2001**, 12, 1411-1415.
931. Hulst R, de Vries NK, Feringa BL. α -Phenylethylamine based chiral phospholidines; new agents for the determination of the enantiomeric excess of chiral alcohols, amines and thiols by means of ^{31}P NMR. *Tetrahedron: Asymmetry* **1994**, 5, 699-708.
932. de Parrodi CA, Moreno GE, Quintero L, Juaristi E. Application of phosphorylated reagents derived from N,N'-di-[(S)- α -phenylethyl]-cyclohexane-1,2-diamines in the determination of the enantiomeric purity of chiral alcohols. *Tetrahedron: Asymmetry* **1998**, 9, 2093-2099.
933. Moreno GE, Mastranzo VM, Quintero L, de Parrodi CA, Juaristi E. The Use of N,N'-di[α -phenylethyl]-diamines as phosphorylated chiral derivatizing agents for the determination of the enantiomeric purity of chiral secondary alcohols. *Rev. Soc. Quim. Mex.* **2003**, 47, 127-129.

934. Devitt PG, Mitchell MC, Weetman JM, Taylor RJ, Kee TP. Accurate in situ $^{31}\text{P}\{^1\text{H}\}$ assay of enantiopurity in α -hydroxyphosphonate esters using a diazaphospholidine derivatizing agent. *Tetrahedron: Asymmetry* **1995**, 6, 2039-2044.
935. Davies SR, Mitchell MC, Cain CP, Devitt PG, Taylor RJ, Kee TP. Phospho-transfer catalysis. On the asymmetric hydrophosphonylation of aldehydes. *J. Organomet. Chem.* **1998**, 550, 29-57.
936. Chauvin A, Bernardinelli G, Alexakis A. Determination of the absolute configuration of chiral aryl-alkyl carbinols using organophosphorus diamine derivatizing agents by ^{31}P NMR spectroscopy. *Tetrahedron: Asymmetry* **2004**, 15, 1857-1879.
937. Reymond S, Brunel JM, Buono G. New chiral organophosphorus derivatizing agent for the determination of enantiomeric composition of chloro- and bromohydrins by ^{31}P NMR spectroscopy. *Tetrahedron: Asymmetry* **2000**, 11, 1273-1278.

P(III) reagents –diol ligands

938. Brunel J, Pardigon O, Maffei M, Buono G. The use of (4R,5R)-dicarboalkoxy 2-chloro 1,3,2-dioxaphospholanes as new chiral derivatizing agents for the determination of enantiomeric purity of alcohols by ^{31}P NMR. *Tetrahedron: Asymmetry* **1992**, 3, 1243-1246.
939. Che C, Zhang ZN, Huang GL, Wang XX, Qin ZH. Using the chiral organophosphorus derivatizing agents for determination of the enantiomeric composition of chiral carboxylic acids by ^{31}P NMR spectroscopy. *Chin. Chem. Lett.* **2004**, 15, 675-678.
940. Bredikhin AA, Bredikhina ZA, Nigmatzyanov FF. On the use of seven-membered phosphorus heterocycles based on 2,2'-dihydroxy-1,1'-binaphthalene and 1,4:3,6-dianhydro-D-mannitol in the ^{31}P NMR analysis of the enantiomeric composition of chiral alcohols. *Russ. Chem. Bull.* **1998**, 47, 411-416.
941. Kato N. Direct chirality determination of secondary carbinol by chirality recognition ability of C_2 symmetry 1,1'-binaphthyl-2,2'-diyl phosphoryl chloride. *J. Am. Chem. Soc.* **1990**, 112, 254-257.
942. Li KY, Zhou ZH, Yeung CH, Tang CC. A chiral phosphorus derivatizing agent for the determination of the enantiomeric excess of chiral alcohols, amines by ^{31}P NMR. *Chin. Chem. Lett.* **2001**, 12, 907-908.
943. Li KY, Zhou ZH, Chan ASC, Tang CC. An efficient chiral phosphorus derivatizing agent for the determination of the enantiomeric excess of chiral alcohols and amines by ^{31}P NMR spectroscopy. *Heteroat. Chem.* **2002**, 13, 93-95.
944. Bredikhin AA, Bredikhin ZA, Gaisina LM, Strunskaya EI, Azancheev NM. Cyclic phosphorochloridites (chloridates) based on chiral butane-2,3-diol and dihydrobenzoin as reagents for the analysis of enantiomeric compositions of alcohols by ^{31}P NMR spectroscopy. *Russ. Chem. Bull.* **2000**, 49, 310-313.
945. Alexakis A, Chauvin A. TADDOL organophosphorus derivatising agents for the determination of the enantiomeric excess of chiral alcohols and carboxylic acids by ^{31}P and ^1H NMR spectroscopy. *Tetrahedron: Asymmetry* **2000**, 11, 4245-4248.
946. Bredikhin AA, Strunskaya EI, Azancheev NM, Bredikhina ZA. (4R,5R)-Bis(N,N-dimethylaminocarbonyl)-2-chloro-1,3,2-dioxaphospholane: a convenient reagent for control of enantiomeric composition of chiral alcohols by ^{31}P NMR spectroscopy. *Russ. Chem. Bull.* **1998**, 47, 174-176.
947. Bredikhina ZA, Novikova VG, Azancheev NM, Bredikhin AA. Regiochemistry of cleavage of monosubstituted oxiranes by phosphorochloridites. Enantiomeric composition of oxiranes by ^{31}P NMR spectral data. *Russ. J. Gen. Chem.* **2002**, 72, 1207-1214.

HELOL Phosphite

957. Weix DJ, Dreher SD, Katz TJ. [5]HELOL phosphite: a helically grooved sensor of remote chirality. *J. Am. Chem. Soc.* **2000**, 122, 10027-10032.

958. Wang DZ, Katz TJ. A [5]HELOL analogue that senses remote chirality in alcohols, phenols, amines, and carboxylic acids. *J. Org. Chem.* **2005**, 70, 8497-8502.

TRISPHAT, BINPHAT, BINTROP

704. Vignon SA, Wong J, Tseng H, Stoddart JF. Helical chirality in donor-acceptor catenanes. *Org. Lett.* **2004**, 6, 1095-1098.

992. Lacour J, Vial L, Bernardinelli G. One-step stereoselective synthesis and NMR chiral shift properties of a novel hexacoordinated phosphorus cation. *Org. Lett.* **2002**, 4, 2309-2312.

993. Lacour J, Hebbe-Viton V. Recent developments in chiral anion mediated asymmetric chemistry. *Chem. Soc. Rev.* **2003**, 32, 373-382

Phosphate configuration

995. Webb MR, Trentham DR. Analysis of chiral inorganic [^{16}O , ^{17}O , ^{18}O]thiophosphate and the stereochemistry of the 3-phosphoglycerate kinase reaction. *J. Biol. Chem.* **1980**, 255, 1775-1779.

996. Buchwald SL, Knowles JR. Determination of the absolute configuration of [^{16}O , ^{17}O , ^{18}O]phosphate monoesters by using ^{31}P NMR. *J. Am. Chem. Soc.* **1980**, 102, 6601-6602.

997. Arnold JRP, Lowe G. Synthesis and stereochemical analysis of chiral inorganic [^{16}O , ^{17}O , ^{18}O]thiophosphate. *J. Chem. Soc., Chem. Commun.* **1986**, 865-867.

998. Bruzik K, Tsai M. Phospholipids chiral at phosphorus. 5. Synthesis and configurational analysis of chiral [^{17}O , ^{18}O]phosphatidylethanolamine. *J. Am. Chem. Soc.* **1984**, 106, 747-754.

999. Lowe G. Chiral [^{16}O , ^{17}O , ^{18}O]phosphate esters. *Acc. Chem. Res.* **1983**, 16, 244-251.

1000. Speckhard DC, Pecoraro VL, Knight WB, Cleland WW. Determination of the absolute configurations of the isomers of triamminecobalt(III) adenosine triphosphate. *J. Am. Chem. Soc.* **1986**, 108, 4167-4171.

1001. Coderre JA, Mehdi S, Demou PC, Weber R, Traficante DD, Gerlt JA. Oxygen chiral phosphodiester. 3. Use of ^{17}O NMR spectroscopy to demonstrate configurational differences in the diastereomers of cyclic 2'-deoxyadenosine 3',5'-[^{17}O , ^{18}O]monophosphate. *J. Am. Chem. Soc.* **1981**, 103, 1870-1872.

1002. Bentrude WG, Sopchik AE, Gajda T. Stereo- and regiochemistries of the oxidations of 2-methoxy-5-tert-butyl-1,3,2-dioxaphosphorinanes and the cyclic methyl 3',5'-phosphite of thymidine by $\text{H}_2\text{O}/\text{I}_2$ and O_2/AIBN to P-chiral phosphates. ^{17}O NMR assignment of phosphorus configuration to the diastereomeric thymidine cyclic methyl 3',5'-monophosphates. *J. Am. Chem. Soc.* **1989**, 111, 3981-3987.

Selenium-based reagent

1006. Silks LA, Dunlap RB, Odom JD. Quantitative detection of remotely disposed chiral centers using selenium-77 nuclear magnetic resonance spectroscopy. *J. Am. Chem. Soc.* **1990**, 112, 4979-4982.

1007. Silks LA, Peng J, Odom JD, Dunlap RB. Synthesis of (4S,5R)-(-)-4-methyl-5-phenyloxazolidine-2-selone: a chiral auxiliary reagent capable of detecting the enantiomers of (R,S)-lipoic acid by ^{77}Se nuclear magnetic resonance spectroscopy. *J. Chem. Soc., Perkin Trans. 1* **1991**, 2495-2498.

1008. Peng J, Odom JD, Dunlap RB, Silks LA. Use of a selone chiral derivatizing agent for the absolute configurational assignment of stereogenic centers. *Tetrahedron: Asymmetry* **1994**, 5, 1627-1630.
1009. Peng J, Barr ME, Ashburn DA, Lebioda L, Garber AR, Martinez RA, Odom JD, Dunlap RB, Silks LA. Synthesis and characterization of acylated chiral oxazolidine-2-selones: selone chiral derivatizing agents for the detection and quantitation of remotely disposed chiral centers. *J. Org. Chem.* **1995**, 60, 5540-5549.
1010. Smith AB, Friestad GK, Barbosa J, Bertounesque E, Duan JJ, Hull KG, Iwashima M, Qiu Y, Spoors PG, Salvatore BA. Total synthesis of (+)-calyculin A and (–)-calyculin B: cyanotetraene construction, asymmetric synthesis of the C(26-37) oxazole, fragment assembly, and final elaboration. *J. Am. Chem. Soc.* **1999**, 121, 10478-10486.
1011. Wu R, Odom JD, Dunlap RB, Silks LA. Simple enantiomeric excess determination of alcohols using chiral selones and ⁷⁷Se NMR spectroscopy. *Tetrahedron: Asymmetry* **1995**, 6, 833-834.
1012. Wu R, Odom JD, Dunlap RB, Silks LA. Reaction of alcohols (via the Mitsunobu reaction) and alkyl halides with chiral selone derivatizing agents. *Tetrahedron: Asymmetry* **1999**, 10, 1465-1470.
1013. Wu R, Hernández G, Odom JD, Dunlap RB, Silks LA. Simple enantiomeric excess determination of amines using chiral selones: unusual N-H···Se bonding detected by HMQC ¹H/⁷⁷Se NMR spectroscopy. *Chem. Commun.* **1996**, 1125-1126.

Cyclodextrins - Native

1048. Greatbanks D, Pickford R. Cyclodextrins as chiral complexing agents in water, and their application to optical purity measurements. *Magn. Reson. Chem.* **1987**, 25, 208-215.
1049. Casy AF, Mercer AD. Application of cyclodextrins to chiral analysis by ¹H NMR spectroscopy. *Magn. Reson. Chem.* **1988**, 26, 765-774.

Cyclodextrins - Permethylated

1071. Botsi A, Yannakopoulou K, Hadjoudis E, Perly B. NMR differentiation of enantiomeric (+)- and (–)- α -pinene via complexation with cyclodextrins in water. *J. Chem. Soc., Chem. Commun.* **1993**, 1085-1086.
1081. Taylor A, Williams DAR, Wilson ID. Derivatized β -cyclodextrins combined with high field NMR for enantiomer analysis: application to ICI 185,282 (5(Z)-7-([2RS,4RS,5SR]-4-o-hydroxyphenyl-2-trifluoromethyl-1,3-dioxan-5-yl) heptenoic acid). *J. Pharm. Biomed. Anal.* **1991**, 9, 493-496.
1082. Kano K, Kato Y, Kodera M. Mechanism for chiral recognition of binaphthyl derivatives by cyclodextrins. *J. Chem. Soc., Perkin Trans. 2*, **1996**, 1211-1217.
1083. Park KK, Park JM. ¹H NMR study of the inclusion complexes of chiral aromatic guests with β -cyclodextrin and its derivatives: discrimination of aromatic protons and chiral recognition. *Bull. Korean Chem. Soc.* **1996**, 17, 1052-1056.
1084. Chankvetadze B, Burjanadze N, Pintore G, Strickmann D, Bergenthal D, Blaschke G. Chiral recognition of verapamil by cyclodextrins studied with capillary electrophoresis, NMR spectroscopy, and electrospray ionization mass spectrometry. *Chirality* **1999**, 11, 635-644.
1086. Uccello-Barretta G, Balzano F, Caporusso AM, Salvadori P. Direct determination of the enantiomeric purity of chiral trisubstituted allenes by using permethylated cyclodextrin as a chiral solvating agent. *J. Org. Chem.* **1994**, 59, 836-839.

1087. Uccello-Barretta G, Balzano F, Caporusso AM, Iodice A, Salvadori P. Permethylated β -cyclodextrin as chiral solvating agent for the NMR assignment of the absolute configuration of chiral trisubstituted allenes. *J. Org. Chem.* **1995**, *60*, 2227-2231.

1088. Uccello-Barretta G, Balzano F, Salvadori P, Lazzaroni R, Caporusso AM, Menicagli R. Different NMR approaches to the chiral analysis of trisubstituted allenes devoid of polar functional groups and aromatic hydrocarbons. *Enantiomer* **1996**, *1*, 365-375.

1089. Uccello-Barretta G, Balzano F, Cuzzola A, Menicagli R, Caporusso AM, Salvadori P. Permethylated cyclodextrins as chiral solvating agents for the determination of the enantiomeric composition of apolar substrates by NMR. *Gazz. Chim. Ital.* **1997**, *127*, 383-386.

Cyclodextrins - Carboxymethylated

1117. Dignam CF, Randall LA, Blacken RD, Cunningham PR, Lester SG, Brown MJ, French SC, Aniagyei SE, Wenzel TJ. Carboxymethylated cyclodextrin derivatives as chiral NMR discriminating agents. *Tetrahedron: Asymmetry* **2006**, *17*, 1199-1208.

1118. Endresz G, Chankvetadze B, Bergenthal D, Blaschke G. Comparative capillary electrophoretic and nuclear magnetic resonance studies of the chiral recognition of racemic metomidate with cyclodextrin hosts. *J. Chromatogr. A* **1996**, *732*, 133-142.

1125. Smith KJ, Wilcox JD, Mirick GE, Wacker LS, Ryan NS, Vensel DA, Readling R, Domush HL, Amonoo EP, Shariff SS, Wenzel TJ. Calix[4]arene, calix[4]resorcarene, and cyclodextrin derivatives and their lanthanide complexes as chiral NMR shift reagents. *Chirality* **2003**, *15*, S150-S158.

1126. Wenzel TJ, Amonoo EP, Shariff SS, Aniagyei SE. Sulfated and carboxymethylated cyclodextrins and their lanthanide complexes as chiral NMR discriminating agents. *Tetrahedron: Asymmetry* **2003**, *14*, 3099-3104.

Crown ether – primary amines

1142. Machida Y, Nishi H, Nakamura K. Nuclear magnetic resonance studies for the chiral recognition of the novel chiral stationary phase derived from 18-crown-6 tetracarboxylic acid. *J. Chromatogr. A* **1998**, *810*, 33-41.

1144. Wenzel TJ, Thurston JE. (+)-(18-Crown-6)-2,3,11,12-tetracarboxylic acid and its ytterbium(III) complex as chiral NMR discriminating agents. *J. Org. Chem.* **2000**, *65*, 1243-1248.

1145. Wenzel TJ, Thurston JE. Enantiomeric discrimination in the NMR spectra of underivatized amino acids and α -methyl amino acids using (+)-(18-crown-6)-2,3,11,12-tetracarboxylic acid. *Tetrahedron Lett.* **2000**, *41*, 3769-3772.

1146. Wenzel TJ, Freeman BE, Sek DC, Zopf JJ, Nakamura T, Yongzhu J, Hirose K, Tobe Y. Chiral recognition in NMR spectroscopy using crown ethers and their ytterbium(III) complexes. *Anal. Bioanal. Chem.* **2004**, *378*, 1536-1547.

1147. Machida Y, Kagawa M, Nishi H. Nuclear magnetic resonance studies for the chiral recognition of (+)-(R)-18-crown-6-tetracarboxylic acid to amino compounds. *J. Pharm. Biomed. Anal.* **2003**, *30*, 1929-1942.

Crown ether – secondary amines

1141. Lovely AE, Wenzel TJ. Chiral NMR discrimination of secondary amines using (18-crown-6)-2,3,11,12-tetracarboxylic acid. *Org. Lett.* **2006**, *8*, 2823-2826.

“Chiral NMR Discrimination of Amines: Analysis of Secondary, Tertiary and Prochiral Amines using (18-Crown-6)-2,3,11,12-tetracarboxylic Acid,” Lovely, A. E., Wenzel, T. J., *Chirality*, in press.

“Chiral NMR Discrimination of Piperidines and Piperazines Using (18-Crown-6)2,3,11,12-tetracarboxylic Acid,” Lovely, A.E.; Wenzel, T.J., *Journal of Organic Chemistry*, **2006**, 71, 9178-9182.

“Chiral NMR Discrimination of Pyrrolidines Using (18-Crown-6)2,3,11,12-tetracarboxylic Acid,” Lovely, A.E.; Wenzel, T.J., *Tetrahedron Asymmetry*, **2006**, 17, 2642-2648.

Calix[4]resorcarenes

1186. Yanagihara R, Tominaga M, Aoyama Y. Chiral host-guest interaction. A water-soluble calix[4]resorcarene having L-proline moieties as a non-lanthanide chiral NMR shift reagent for chiral aromatic guests in water. *J. Org. Chem.* **1994**, 59, 6865-6867.

1187. Dignam CF, Richards CJ, Zopf JJ, Wacker LS, Wenzel TJ. An enantioselective NMR shift reagent for cationic aromatics. *Org. Lett.* **2005**, 7, 1773-1776.

1188. Dignam CF, Zopf JJ, Richards CJ, Wenzel TJ. Water-soluble calix[4]resorcarenes as enantioselective NMR shift reagents for aromatic compounds. *J. Org. Chem.* **2005**, 70, 8071-8078.

Lanthanide complexes

1248. Whitesides GM, Lewis DW. Tris[3-(tert-butylhydroxymethylene)-d-camphorato]europium(III). A reagent for determining enantiomeric purity. *J. Am. Chem. Soc.* **1970**, 92, 6979-6980.

1249. Goering HL, Eikenberry JN, Koerner GS. Tris[3-(trifluoromethylhydroxymethylene)-d-camphorato]europium(III). A chiral shift reagent for direct determination of enantiomeric compositions. *J. Am. Chem. Soc.* **1971**, 93, 5913-5914.

1250. Fraser RR, Petit MA, Saunders JK. Determination of enantiomeric purity by an optically active nuclear magnetic resonance shift reagent of wide applicability. *Chem. Commun.* **1971**, 1450-1451.

1251. Wenzel TJ. *NMR Shift Reagents*. CRC Press, Inc: Boca Raton, FL, **1978**.

1252. McCreary MD, Lewis DW, Wernick DL, Whitesides GM. The determination of enantiomeric purity using chiral lanthanide shift reagents. *J. Am. Chem. Soc.* **1974**, 96, 1038-1054.

1253. Goering HE, Eikenberry JN, Koerner GS, Lattimer CJ. Direct determination of enantiomeric compositions with optically active nuclear magnetic resonance lanthanide shift reagents. *J. Am. Chem. Soc.* **1974**, 96, 1493-1501.

1314. Bartik K, El Haouaj M, Luhmer M, Collet A, Reisse J. Can monoatomic xenon become chiral? *ChemPhysChem* **2000**, 1, 221-224.

Binuclear lanthanide-silver complexes

1246. Wenzel TJ, Bettis TC, Sadlowski JE, Sievers RE. New binuclear lanthanide NMR shift reagents effective for aromatic compounds. *J. Am. Chem. Soc.* **1980**, 102, 5903-5904.

1247. Wenzel TJ, Sievers RE. Binuclear complexes of lanthanide(III) and silver(I) and their function as shift reagents for olefins, aromatics, and halogenated compounds. *Anal. Chem.* **1981**, 53, 393-399.

1361. Offermann W, Mannschreck A. Chiral recognition of alkene and arene hydrocarbons by ^1H and ^{13}C NMR. Determination of enantiomeric purity. *Tetrahedron Lett.* **1981**, 22, 3227-3230.

1362. Wenzel TJ, Sievers RE. Nuclear magnetic resonance studies of terpenes with chiral and achiral lanthanide(III)-silver(I) binuclear shift reagents. *J. Am. Chem. Soc.* **1982**, 104, 382-388.

1363. Wenzel TJ, Lalonde DR. New binuclear NMR shift reagents for olefins and aromatics. *J. Org. Chem.* **1983**, 48, 1951-1954.
1364. Offermann W, Mannschreck A. Application of NMR spectroscopy of chiral association complexes. 14. Chiral recognition of terpene and cyclohexene hydrocarbons by ^1H and ^{13}C NMR. *Org. Magn. Reson.* **1984**, 22, 355-363.
1365. Krasutskii PA, Rodionov VN, Tikhonov VP, Yurchenko AG. Silver d-camphor-10-sulfonate as a new chiral reagent for enantiomeric resolution of olefins. *Teor. Eksp. Khim.* **1984**, 20, 58-63.
1366. Wenzel TJ, Ruggles AC, Lalonde DR. Binuclear lanthanide(III)—silver(I) NMR shift reagents: investigations of new achiral and chiral analogs. *Magn. Reson. Chem.* **1985**, 23, 778-783.
1367. Baldovini N, Tomi F, Casanova J. Enantiomeric differentiation of terpenic olefins by carbon-13 NMR using chiral binuclear shift reagents. *Magn. Reson. Chem.* **2001**, 39, 621-624.
1368. Peterson PE, Jensen BL. The preparation of 1,3-bis(trimethylsilyl)allene and the observation of the enantiomers using olefin complexing chiral shift reagents. *Tetrahedron Lett.* **1984**, 25, 5711-5714.
1369. Mannschreck A, Munninger W, Burgemeister T, Gore J, Cazes B. Chiral recognition of allenic hydrocarbons by ^1H NMR. *Tetrahedron* **1986**, 42, 399-408.
1370. Pasto DJ, Sugi KD. On the maximum rotation and the solvobromination and –mercuration of enantioenriched 1,3-dimethylallene. *J. Org. Chem.* **1991**, 56, 4157-4160.
1373. Pasto DJ, Sugi KD. Stereochemical features of the (2 + 2) cycloaddition reactions of chiral allenes. 3. The cycloaddition of enantioenriched 1,3-dimethylallene with the symmetrically 1,1-disubstituted alkenes 1,1-dichloro-2,2-difluoroethene and 1,1-diphenylethene. *J. Org. Chem.* **1992**, 57, 12-17.
1379. Wenzel TJ, Zaia J. Lanthanide tetrakis (β -diketonates) as effective NMR shift reagents for organic salts. *J. Org. Chem.* **1985**, 50, 1322-1324.
1380. Wenzel TJ, Zaia J. Organic-soluble lanthanide nuclear magnetic resonance shift reagents for sulfonium and isothiuronium salts. *Anal. Chem.* **1987**, 59, 562-567.
1381. Green TK, Whetstone JR, Son E. Enantiomeric purity of alkylmethylphenylsulfonium ions with chiral NMR shift reagents: racemization by pyramidal inversion as observed by ^1H NMR spectroscopy. *Tetrahedron: Asymmetry* **1997**, 8, 3175-3181.

Reduction of broadening – lanthanide complexes

1255. Fraser RR, Stothers JB, Tan CT. Determination of optical purity with chiral shift reagents and ^{13}C magnetic resonance. *J. Magn. Reson.* **1973**, 10, 95-97.
1256. Pearce GT, Gore WE, Silverstein RM. Synthesis and absolute configuration of multistriatin. *J. Org. Chem.* **1976**, 41, 2797-2803.
1257. Belleney J, Bui C, Carrière FJ. Determination of the enantiomeric purity of disubstituted β -lactams by NMR spectroscopy using a chiral shift reagent. *Magn. Reson. Chem.* **1990**, 28, 606-611.
1258. Omata K, Aoyagi S, Kabuto K. Observing the enantiomeric ^1H chemical shift non-equivalence of several α -amino ester signals using tris[3-(trifluoromethylhydroxy-methylene)-(+)-camphorato]samarium(III): a chiral lanthanide shift reagent that causes minimal line bordering. *Tetrahedron: Asymmetry* **2004**, 15, 2351-2356.

1259. Sweeting LM, Crans DC, Whitesides GM. Determination of enantiomeric purity of polar substrates with chiral lanthanide NMR shift reagents in polar solvents. *J. Org. Chem.* **1987**, 52, 2273-2276.
1260. Rothchild R, Sanders K, Venkatasubban KS. NMR studies of drugs. Use of lanthanide shift reagents in polar solvent with thalidomide. *Spectrosc. Lett.* **1993**, 26, 597-619.
1261. DeArment M, Eastabrooks M, Venkatasubban KS, Benschafut R, Rothchild R, Wyss H. NMR studies of drugs. Application of a chiral lanthanide shift reagent to methoxamine in chloroform-d or acetonitrile-d₃. *Spectrosc. Lett.* **1994**, 27, 533-555.
1262. Polívka Z, Buděšínský M, Holubek J, Schneider B, Šedivý Z, Svátek E, Matoušová O, Metyš J, Valchář M, Souček R, Protiva M. 4H-Benzo[4,5]cyclohepta[1,2-b]thiophenes and 9,10-dihydro derivatives - sulfonium analogues of pizotifen and ketotifen; chirality of ketotifen; synthesis of the 2-bromo derivative of ketotifen. *Collect. Czech. Chem. Commun.* **1989**, 54, 2443-2469.
1263. Peterson PE, Stepanian M. Methodology for the analysis of products from asymmetric syntheses using chiral NMR shift reagents. Relative complexation constants of enantiomers. *J. Org. Chem.* **1988**, 53, 1907-1911.

Assignment of absolute configuration – lanthanide complexes

1340. Červinka O, Maloň P, Trška P. Asymmetric reactions. XLVII. NMR-shift differences of enantiotopic proton signals in some alkylaryl methanols induced by tris[3-trifluoroacetyl-(+)-camphorato]europium(III). *Collection Czechoslov. Chem. Commun.* **1973**, 38, 3299-3301.
1341. Sullivan GR, Ciavarella D, Mosher HS. Correlation of configuration of chiral secondary carbinols by use of a chiral lanthanide nuclear magnetic resonance shift reagent. *J. Org. Chem.* **1974**, 39, 2411-2412.
1342. Yoshida N, Morita H, Oyama K, Lee H-H. Enzymatic syntheses of chiral sec-alcohols and their NMR chemical shifts induced by a chiral lanthanide shift reagent. *Chem. Express* **1989**, 4, 721-724.
1343. Ghosh I, Zeng H, Kishi Y. Application of chiral lanthanide shift reagents for assignment of absolute configuration of alcohols. *Org. Lett.* **2004**, 6, 4715-4718.
1344. Ghosh I, Kishi Y, Tomoda H, Omura S. Use of a chiral praseodymium shift reagent in predicting the complete stereostructure of glisoprenin A. *Org. Lett.* **2004**, 6, 4719-4722.
1345. Adams CM, Ghosh I, Kishi Y. Validation of lanthanide chiral shift reagents for determination of absolute configuration: total synthesis of glisoprenin A. *Org. Lett.* **2004**, 6, 4723-4726.
1346. Capillon J, Lacombe L. Effets des substituants dans les spectres de rmn de benzhydrols substitués en présence de Eu (dcm)₃. *Can. J. Chem.* **1979**, 57, 1446-1450.
1347. Mioskowski C, Solladie G. Enantiomeric purity and absolute configuration of β-hydroxy- β-(trifluoromethyl)-β-phenylpropionic acid obtained by asymmetric synthesis. *Tetrahedron* **1973**, 29, 3669-3674.
1348. Azzolina O, Ghislandi V. Antiphlogistic aryloxypropionic acids: configurational study. *Farmaco* **1993**, 48, 713-724.
1349. Ajisaka K, Kamisaku M, Kainosho M. Enantiomeric shift difference induced by the lanthanide shift reagent: its correlation with absolute configuration of α-amino acids. *Chem. Lett.* **1972**, 857-858.
1350. Cativiala C, Diaz de Villegas MD, Garcia JI, Mayoral JA, Meléndez E. Determination of the enantiomeric purity by ¹H NMR with Eu(tfc)₃ of β-hetarylalanine derivatives. Correlation of the enantiomeric shift difference with absolute configuration. *Bull. Soc. Chim. Belg.* **1984**, 93, 479-482.

1351. Bus J, Lok CM, Groenewegen A. Determination of enantiomeric purity of glycerides with a chiral PMR shift reagent. *Chem. Phys. Lipids* **1976**, 16, 123-132.
1352. Hudhomme P, Duguay G. Control of the enantiomeric purity and correlation with the absolute configuration of N-protected 2-cyanoglycinates. *Tetrahedron: Asymmetry* **1993**, 4, 1891-1900.
1353. Jankowski K, Rabcenko A. Application of an optically active nuclear magnetic resonance shift reagent to configurational problems. *J. Org. Chem.* **1975**, 960-961.
1354. White JD, Somers TC, Reddy GN. Degradation and absolute configurational assignment to C₃₄-botryococcene. *J. Org. Chem.* **1992**, 57, 4991-4998.
1355. White JD, Somers TC, Reddy GN. Absolute configuration of (–)-botryococcene. *J. Am. Chem. Soc.* **1986**, 108, 5352-5353.
1356. Nishida R, Kuwahara Y, Fukami H, Ishii S. Female sex pheromone of the German cockroach, *Blattella germanica* (L.) (Orthoptera: Blattellidae), responsible for male wing-raising. IV. The absolute configuration of the pheromone, 3,11-dimethyl-2-nonacosanone. *J. Chem. Ecol.* **1979**, 5, 289-297.
1357. Yeh HJC, Balani SK, Yagi H, Greene RME, Sharma ND, Boyd DR, Jerina DM. Use of chiral lanthanide shift reagents in the determination of enantiomer composition and absolute configuration of epoxides and arene oxides. *J. Org. Chem.* **1986**, 51, 5439-5443.
1358. Langin-Lantéri M, Fonbonne C, Huet J, Petit-Ramel M. Chiral phenyl-substituted chalcone epoxides. I. Correlative determination of absolute configuration by ¹H NMR lanthanide-induced shifts. *Magn. Reson. Chem.* **1987**, 25, 216-218.

Water-soluble lanthanide complexes

1386. Kabuto K, Sasaki Y. The europium(III)-(R)-propylenediaminetetra-acetate ion: a promising chiral shift reagent for ¹H N.M.R. spectroscopy in aqueous solution. *J. Chem. Soc., Chem. Commun.* **1984**, 316-318.
1387. Kabuto K, Sasaki Y. Highly consistent correlation between absolute configuration of α -amino acids and their shift induced by the N.M.R. chiral shift reagent propylenediaminetetraacetatoeuropium(III) in aqueous solution. *J. Chem. Soc., Chem. Commun.* **1987**, 670-671.
1388. Kabuto K, Sasaki Y. Consistent correlation between absolute configuration of α -hydroxy carboxylic acids and lanthanoid induced shift by an NMR chiral shift reagent in aqueous solution. *Chem. Lett.* **1989**, 385-388.
1389. Kabuto K, Sasaki Y. A facile NMR method for assigning absolute configuration of underivatized α -methyl- α -amino acids using a chiral lanthanoid shift reagent for aqueous solution. *Tetrahedron Lett.* **1990**, 31, 1031-1034.
1390. Kabuto K, Sasaki K, Sasaki Y. Absolute configuration of aldonic acids and lanthanoid induced shift by the chiral shift reagent propylenediaminetetraacetatoeuropium(III) in aqueous solution. *Tetrahedron: Asymmetry* **1992**, 3, 1357-1360.
1391. Sasaki M, Omata K, Kabuto K, Sasaki Y. Enantiomer signal separation of β -amino acids induced by Eu(III)-pdta; relation between the absolute configurations and relative shifts of enantiomer signals. *Kidorui* **2003**, 42, 198-199.
1393. Ogasawara K, Omata K, Kabuto K, Jin H, Sasaki Y. Optically active Ce(III) - propylenediaminetetracetate complex: a chiral shift reagent for aqueous solution causing less signal broadening. *Kidorui* **1999**, 34, 152-153.
1394. Inamoto A, Ogasawara K, Omata K, Kabuto K, Sasaki Y. Samarium(III)-propylenediaminetetraacetate complex: a water-soluble chiral shift reagent for use in high-field NMR. *Org. Lett.* **2000**, 2, 3543-3545.

1395. Kabuto K, Hazama R, Umakoshi K, Sasaki Y. A new lanthanide complex, [EuIII((R)-tppn)Cl₂](ClO₄), as a chiral NMR shift reagent for aqueous solution. *Kidorui* **1995**, 26, 244-245.
1396. Hazama R, Umakoshi K, Kabuto C, Kabuto K, Sasaki Y. A europium(III)-N,N,N',N'-tetrakis(2-pyridylmethyl)-(R)-propylenediamine complex as a new chiral lanthanide NMR shift reagent for aqueous neutral solution. *Chem. Commun.* **1996**, 15-16.
1397. Sato J, Omata K, Kabuto K, Jin H, Umakoshi K, Sasaki Y. Properties of chiral Ce(III)-tppn complex as a chiral shift reagent in aqueous solution. *Kidorui*, **1998**, 32, 58-59.
1398. Sato J, Jin H, Omata K, Kabuto K, Sasaki Y. Resolution of enantiomer signals by diamagnetic lanthanum(III)-N,N,N',N'-tetrakis(2-pyridinylmethyl)-(R)-propylenediamine complex in ¹H NMR. *Enantiomer*, **1999**, 4, 147-150.

Palladium dimers

1411. Hockless DCR, Gugger PA, Leung P, Mayadunne RC, Pabel M, Wild SB. Facile interconversions between diastereomers of chloro-bridged palladium(II) dimers of orthometallated (±)-dimethyl[1-(1-naphthyl)ethyl]amine. *Tetrahedron* **1997**, 53, 4083-4094.
1413. Wild SB. Resolutions of tertiary phosphines and arsines with orthometallated palladium(II)-amine complexes. *Coord. Chem. Rev.* **1997**, 166, 291-311.
1415. Kyba EP, Rines SP. A spectroscopic method for the determination of optical purities of chiral, chelating diphosphines. *J. Org. Chem.* **1982**, 47, 4800-4801.
1416. Roberts NK, Wild SB. Resolutions using metal complexes. Synthesis, separation into diastereoisomers, and resolution of o-phenylenebis(methylphenylphosphine) using palladium complexes containing optically active orthometallated dimethyl(α-methylbenzyl)amines. *J. Am. Chem. Soc.* **1979**, 101, 6254-6260.
1417. Kashiwabara K, Kinoshita I, Fujita J. Optical resolution of 2-aminoethyl-n-butylphenylphosphine and the bis(acetylacetonato) cobalt(III) complex containing the resolved aminophosphine. *Chem. Lett.* **1978**, 673-676.
1418. Kyba EP, Rines SP. A spectroscopic method for the determination of optical purities of chiral, chelating diphosphines. *J. Org. Chem.* **1982**, 47, 4800-4801.
1419. Allen DG, McLaughlin GM, Robertson GB, Steffen WL, Salem G, Wild SB. Resolutions involving metal complexation. Preparation and resolution of (R,S)-methylphenyl(8-quinolyl)phosphine and its arsenic analogue. Crystal and molecular structure of (+)₅₈₉-(R)-dimethyl(1-ethyl-α-naphthyl)aminato-C²,N]-[(S)-methylphenyl(8-quinolyl)phosphine]palladium(II) hexafluorophosphate. *Inorg. Chem.* **1982**, 21, 1007-1014.
1420. Martin JWL, Palmer JAL, Wild SB. Resolutions involving metal complexation. Resolution of (±)-1-amino-2-(methylphenylarsino)ethane and its phosphorus analogue. Stereochemistry and stability of square-planar bis(bidentate) complexes of bivalent palladium and platinum. *Inorg. Chem.* **1984**, 23, 2664-2668.
1427. Albert J, Bosque R, Cadena JM, Granell JR, Muller G, Ordinas JI. Synthesis and resolution of benzylisopropylphenylphosphine, a monodentate P-chiral ligand, *Tetrahedron: Asymmetry* **2000**, 11, 3335-3343.
1428. Albert J, Bosque R, Cadena JM, Delgado S, Granell J, Muller G, Ordinas JI, Bardia MF, Solans X. Synthesis, resolution, and reactivity of benzylmesitylphenylphosphine: a new P-chiral bulky ligand. *Chem. Eur. J.* **2002**, 8, 2249-2287.
1431. Dunina VV, Golovan EB, Gulyukina NS, Buyevich AV. Enantiomeric discrimination in the complexation of ortho-palladated α-arylalkylamines with the racemic tert-butylmethylphenylphosphine. *Tetrahedron: Asymmetry* **1995**, 6, 2731-2746.

1432. Dunina VV, Kuz'mina LG, Kazakova MY, Grishin YK, Veits YA, Kazakova EI. Ortho-palladated α -phenylalkylamines for enantiomeric purity determination of monodentate P*-chiral phosphines. *Tetrahedron: Asymmetry* **1997**, 8, 2537-2545.

1433. Dunina VV, Kuz'mina LG, Rubina MY, Grishin YK, Veits YA, Kazakova EI. A resolution of the monodentate P*-chiral phosphine $\text{P}^+\text{Bu}^1\text{C}^6\text{H}_4\text{Br}-4$ and its NMR-deduced absolute configuration. *Tetrahedron: Asymmetry* **1999**, 10, 1483-1497.

Palladium/platinum for olefins

1442. Parker D, Taylor RJ. Simple organometallic chiral derivatising agents for the ^{31}P N.M.R.s. assay of the enantiomeric purity of certain η^2 -donors. *J. Chem. Soc., Chem. Commun.* **1987**, 1781-1783.

1443. Parker D, Taylor RJ. Scope and limitations of organometallic chiral derivatising agents for the ^{31}P NMR determination of the enantiomeric purity of chiral η^2 donors. *Tetrahedron* **1988**, 44, 2241-2248.

1450. Fulwood R, Parker D, Ferguson G, Kaltner B. Crystal structure and further applications of the chiral derivatising agent η^2 -ethene platinum-DIOP. *J. Organomet. Chem.* **1991**, 419, 269-276.

Platinum-amine complexes

1087. Uccello-Barretta G, Balzano F, Caporusso AM, Iodice A, Salvadori P. Permethylated β -cyclodextrin as chiral solvating agent for the NMR assignment of the absolute configuration of chiral trisubstituted allenes. *J. Org. Chem.* **1995**, 60, 2227-2231.

1453. Salvadori P, Uccello-Barretta G, Bertozzi S, Settambolo R, Lazzaroni R. Determination of the enantiomeric purity of chiral allyl alcohols and allyl ethers by ^{195}Pt NMR Spectroscopy. *J. Org. Chem.* **1988**, 53, 5678-5770.

1454. Salvadori P, Uccello-Barretta G, Lazzaroni R, Caporusso AM. New method for the enantiomeric excess determination of chiral trisubstituted allenes by ^{195}Pt NMR of trans-dichloro[(S)- α -methylbenzylamine](allene)platinum(II) complexes. *J. Chem. Soc., Chem. Commun.* **1990**, 1121-1123.

1455. Uccello-Barretta G, Bernardini R, Balzano F, Lazzaroni R, Salvadori P. On the high efficiency of cis-dichloro[(S)- α -methylbenzylamine](ethylene)platinum(II) as chiral derivatizing agent for the determination of the enantiomeric composition of chiral unsaturated ethers by ^{195}Pt -NMR spectroscopy: a spectroscopic conformational and configurational characterization in solution of diastereoisomeric complexes cis-dichloro[(S)- α -methylbenzylamine][(S)- and (R)-3-phenyl-3-methoxybut-1-ene]platinum(II). *J. Organomet. Chem.* **2000**, 605, 68-73.

1456. Uccello-Barretta G, Bernardini R, Lazzaroni R, Salvadori P. Use of cis-dichloro[(S)- α -methylbenzylamine](ethylene)Pt(II) as a chiral derivatizing agent for the determination by ^{195}Pt -NMR of the enantiomeric composition of unsaturated ethers or alcohols having a quaternary chiral carbon atom. *J. Organomet. Chem.* **2000**, 598, 174-178.

1457. Uccello-Barretta G, Bernardini R, Lazzaroni R, Salvador P. $[\text{PtCl}_3(\text{C}_2\text{H}_4)]^-[(\text{S},\text{S})-(\text{PhMeCH})_2\text{NH}_2]^+$: a new and versatile ionic organometallic chiral derivatizing agent for the determination of the enantiomeric composition of chiral unsaturated compounds by ^{195}Pt NMR spectroscopy. *Org. Lett.* **2000**, 2, 1795-1798.

1458. Uccello-Barretta G, Bernardini R, Balzano F, Venturelli F, Caporusso AM, Salvadori P. trans- $[\text{PtCl}_2(\text{Am})(\text{C}_2\text{H}_4)]$ and $[\text{PtCl}_3(\text{C}_2\text{H}_4)]^-[\text{AmH}]^+$, containing binaphthyl secondary amines. Efficient chiral derivatizing agents for the enantiodiscrimination of chiral olefins by ^{195}Pt NMR spectroscopy. *Eur. J. Org. Chem.* **2001**, 3651-3655.

1459. Uccello-Barretta G, Bernardini R, Balzano F, Caporusso AM, Salvadori P. ^{195}Pt NMR determination of the enantiomeric purity and absolute configuration of trisubstituted allenes by using $[\text{PtCl}_3(\text{C}_2\text{H}_4)]^+[(\text{S,S})\text{-(1-NpMeCH)}_2\text{NH}_2]^+$ as CDA. *Org. Lett.* **2001**, 3, 205-207.

1460. Uccello-Barretta G, Bernardini R, Balzano F, Salvadori P. $[\text{PtCl}_3(\text{C}_2\text{H}_4)]^+[\text{AmH}]^+$ complexes containing chiral secondary amines: use as chiral derivatizing agents for the enantiodiscrimination of unsaturated compounds by ^{195}Pt NMR spectroscopy and NMR stereochemical investigation. *J. Org. Chem.* **2001**, 66, 123-129.

1461. Uccello-Barretta G, Bernardini R, Balzano F, Salvadori P. Overall view of the use of chiral platinum(II) complexes as chiral derivatizing agents (CDAs) for the enantiodiscrimination of unsaturated compounds by ^{195}Pt NMR. *Chirality* **2002**, 14, 484-489.

Rhodium dimer

1463. Duddeck H. $\text{Rh}_2[\text{MTPA}]_4$, a dirhodium complex as NMR auxiliary for chiral recognition. *The Chemical Record* **2005**, 5, 1-14.

Zinc tweezer

1522. Ema T, Ouchi N, Doi T, Korenaga T, Sakai T. Highly sensitive chiral shift reagent bearing two zinc porphyrins. *Org. Lett.* **2005**, 7, 3985-3988.

Database methods

74. Lee J, Kobayashi Y, Tezuka K, Kishi Y. Toward creation of a universal NMR Database for the stereochemical assignment of acyclic compounds: proof of concept. *Org. Lett.* **1999**, 1, 2181-2184.

75. MacMillan JB, Molinski TF. Caylobolide A, a unique 36-membered macrolactone from a Bahamian *Lyngbya majuscula*. *Org. Lett.* **2002**, 4, 1535-1538.

141. Kobayashi Y, Lee J, Tezuka K, Kishi Y. Toward creation of a universal NMR database for the stereochemical assignment of acyclic compounds: the case of two contiguous propionate units. *Org. Lett.* **1999**, 1, 2177-2180.

142. Kobayashi Y, Tan C, Kishi Y. Stereochemical assignment of the C21-C38 portion of the desertomycin/oasomycin class of natural products by using universal NMR databases: prediction. *Angew. Chem., Int. Ed.* **2000**, 39, 4279-4281.

143. Tan C, Kobayashi Y, Kishi Y. Stereochemical assignment of the C21-C38 portion of the desertomycin/oasomycin class of natural products by using universal NMR databases: proof. *Angew. Chem., Int. Ed.* **2000**, 39, 4282-4284.

144. Kobayashi Y, Tan C, Kishi Y. Toward creation of a universal NMR database for stereochemical assignment: the case of 1,3,5-trisubstituted acyclic systems. *Helv. Chim. Acta* **2000**, 83, 2562-2571.

145. Kobayashi Y, Tan C, Kishi Y. Toward creation of a universal NMR database for stereochemical assignment: complete structure of the desertomycin/oasomycin class of natural products. *J. Am. Chem. Soc.* **2001**, 123, 2076-2078.

744. Kobayashi Y, Hayashi N, Tan C, Kishi Y. Toward the creation of NMR databases in chiral solvents for assignments of relative and absolute stereochemistry: proof of concept. *Org. Lett.* **2001**, 3, 2245-2248.

745. Hayashi N, Kobayashi Y, Kishi Y. Toward the creation of NMR databases in chiral solvents for assignments of relative and absolute stereochemistry: scope and limitation. *Org. Lett.* **2001**, 3, 2249-2252.

746. Fidanze S, Song F, Szlosek-Pinaud M, Small PLC, Kishi Y. Complete structure of the mycolactones. *J. Am. Chem. Soc.* **2001**, *123*, 10117-10118.
747. Kobayashi Y, Hayashi N, Kishi Y. Toward the creation of NMR databases in chiral solvents for assignments of relative and absolute stereochemistry: NMR desymmetrization of meso compounds. *Org. Lett.* **2001**, *3*, 2253-2255.
748. Kobayashi Y, Hayashi N, Kishi Y. Toward the creation of NMR databases in chiral solvents: bidentate chiral NMR solvents for assignment of the absolute configuration of acyclic secondary alcohols. *Org. Lett.* **2002**, *4*, 411-414.
749. Higashibayashi S, Kishi Y. Assignment of the relative and absolute configurations of acyclic secondary 1,2-diols. *Tetrahedron* **2004**, *60*, 11977-11982.
750. Kobayashi Y, Hayashi N, Kishi Y. Application of chiral bidentate NMR solvents for assignment of the absolute configuration of alcohols: scope and limitation. *Tetrahedron Lett.* **2003**, *44*, 7489-7491.
751. Kobayashi Y, Czechtizky W, Kishi Y. Complete stereochemistry of tetrafibricin. *Org. Lett.* **2003**, *5*, 93-96.
752. Dambruoso P, Bassarello C, Bifulco G, Appendino G, Battaglia A, Fontana G, Gomez-Paloma L. Advances in the universal NMR database approach. 2'-substituted taxanes as probes for an improved protocol of diastereomeric differentiation. *Org. Lett.* **2005**, *7*, 983-986.

Liquid crystals

1548. Meddour A, Canet I, Loewenstein A, Péchiné JM, Courtieu J. Observation of enantiomers, chiral by virtue of isotopic substitution, through deuterium NMR in a polypeptide liquid crystal. *J. Am. Chem. Soc.* **1994**, *116*, 9652-9656.
1557. Canet I, Meddour A, Courtieu J, Canet JL, Salaün J. New, and accurate method to determine the enantiomeric purity of amino acids based on deuterium NMR in a cholesteric lyotropic liquid crystal. *J. Am. Chem. Soc.* **1994**, *116*, 2155-2156.
1558. Chevtchouk T, Ollivier J, Salaün J, Merlet D, Courtieu J. Determination of the region- and enantioselectivity of the enzymatic hydrolyses of succinates based on H,C'COLOC analysis and deuterium NMR in a chiral liquid crystal. *Tetrahedron: Asymmetry* **1997**, *8*, 999-1003.
1559. Canet J, Canet I, Courtieu J, Da Silva S, Gelas J, Troin Y. Acetyl-d₃ chloride: a convenient nonchiral derivatizing agent (NCDA) for a facile enantiomeric excess determination of amines through deuterium NMR. *J. Org. Chem.* **1996**, *61*, 9035-9037.
1560. Chalard P, Bertrand M, Canet I, Théry V, Remuson R, Jeminet G. Determination of absolute configurations of amines and amino acids using nonchiral derivatizing agents (NCDA) and deuterium NMR. *Org. Lett.* **2000**, *2*, 2431-2432.
1561. Meddour A, Loewenstein A, Péchiné J, Courtieu J. An achiral deuterated derivatizing agent for enantiomeric analysis through NMR in a liquid crystalline solvent. *Tetrahedron: Asymmetry* **1997**, *8*, 485-494.
1562. Meddour A, Courtieu J. Achiral deuterated derivatizing agent for enantiomeric analysis of carboxylic acids by NMR in a chiral liquid crystalline solvent. *Tetrahedron: Asymmetry* **2000**, *11*, 3635-3644.
1563. Meddour A, Atkinson D, Loewenstein A, Courtieu J. Enantiomeric analysis of homologous series of secondary alcohols by deuterium NMR spectroscopy in a chiral nematic liquid crystal: influence of molecular geometry on chiral discrimination. *Chem. Eur. J.* **1988**, *4*, 1142-1147.

1564. Tavasli M, Courtieu J, Goss RJM, Meddour A, O'Hagan D. Extreme enantiomeric discrimination of fluoroalkanes using deuterium NMR in chiral liquid crystalline media. *Chem. Commun.* **2002**, 844-845.
1565. Meddour A, Canlet C, Blanco L, Courtieu J. Diastereomeric shape recognition using NMR spectroscopy in a chiral liquid crystalline solvent. *Angew. Chem., Int. Ed.* **1999**, 38, 2391-2393.
1566. Canlet C, Merlet D, Lesot P, Meddour A, Lowenstein A, Courtieu J. Deuterium NMR stereochemical analysis of threo-erythro isomers bearing remote stereogenic centers in racemic and non-racemic liquid crystalline solvents. *Tetrahedron: Asymmetry* **2000**, 11, 1911-1918.
1567. Emsley JW, Lesot P, Courtieu J, Merlet D. The effect of a chiral nematic solvent on the orientational order and conformational distribution of a flexible prochiral solute. *Phys. Chem. Chem. Phys.* **2004**, 6, 5331-5337.
1568. Sarfati M, Aroulanda C, Courtieu J, Lesot P. Enantiomeric recognition of chiral invertomers through NMR in chiral oriented phases: a study of cis-decalin. *Tetrahedron: Asymmetry* **2001**, 12, 737-744.
1569. Lesot P, Lafon O, Kagan HB, Fan C. Study of molecular rotational isomerism using deuterium NMR in chiral oriented solvents. *Chem. Commun.* **2006**, 389-391.
1570. Lesot P, Merlet D, Sarfati M, Courtieu J, Zimmermann H, Luz Z. Enantiomeric and enantiotopic analysis of cone-shaped compounds with C_3 and C_{3v} symmetry using NMR spectroscopy in chiral anisotropic solvents. *J. Am. Chem. Soc.* **2002**, 124, 10071-10082.
1571. Merlet D, Emsley JW, Lesot P, Courtieu J. The relationship between molecular symmetry and second-rank orientational order parameters for molecules in chiral liquid crystalline solvents. *J. Chem. Phys.* **1999**, 111, 6890-6896.
1572. Lafontaine E, Pechine JM, Courtieu J, Mayne CL. Visualization of enantiomers in cholesteric solvents through deuterium NMR. *Liq. Cryst.* **1990**, 7, 293-298.
1573. Bayle J, Courtieu J, Gabetty E, Loewenstein A, Péchiné JM. Enantiomeric analysis in a polypeptide lyotropic liquid crystal through proton decoupled deuterium NMR. *New J. Chem.* **1992**, 16, 837-838.
1574. Canet J, Fadel A, Salaün J, Canet-Fresse I, Courtieu J. Enantiomeric excess analysis of sesquiterpene precursors through proton decoupled deuterium NMR in cholesteric lyotropic liquid crystal. *Tetrahedron: Asymmetry* **1993**, 4, 31-34.
1575. Canet I, Courtieu J, Dauphin G, Gourcy J, Veschambre H. Enantiomeric excess analysis of (2R,3S)-3-deuterio-2-methylcyclohexanone and (1S,2R,3S)-3-deuterio-2-methylcyclohexanol, through deuterium NMR in a polypeptide lyotropic liquid crystal. *Tetrahedron: Asymmetry* **1995**, 6, 333-336.
1576. Canet I, Courtieu J, Loewenstein A, Meddour A, Pechine JM. Enantiomeric analysis in a polypeptide lyotropic liquid crystal by deuterium NMR. *J. Am. Chem. Soc.* **1995**, 117, 6520-6526.
1577. Smadja W, Auffret S, Berdagué P, Merlet D, Canlet C, Courtieu J, Legros J, Boutros A, Fiaud J. Visualisation of axial chirality using $^2\text{H}\{-^1\text{H}\}$ NMR in poly(γ -benzyl L-glutamate), a chiral liquid crystal solvent. *Chem. Commun.* **1997**, 2031-2032.
1578. Canet J, Canet I, Gelas J, Ripoché I, Troin Y. Facile and performant enantiomeric excess analysis of diene iron tricarbonyl complexes through deuterium NMR. *Tetrahedron: Asymmetry* **1997**, 8, 2447-2451.
1579. Villar H, Guibé F, Aroulanda C, Lesot P. Investigation of SmI_2 -mediated cyclisation of δ -iodo- α,β -unsaturated esters by deuterium 2D NMR in oriented solvents. *Tetrahedron: Asymmetry* **2002**, 13, 1465-1475.

1580. Gouriou L, Lloyd-Jones GC, Vyskočil Š, Kočovský P. ^2H -quadrupolar coupling-based analysis of stereochemical and regiochemical memory in the Pd-catalysed allylic alkylation of iso-cinnamyl type substrates employing the chiral monophosphine ligands 'MOP' and 'MAP'. *J. Organomet. Chem.* **2003**, 687, 525-537.
1581. Phillips AR, Sharman GJ. The measurement of high enantiomeric excesses in chiral liquid crystals using ^{19}F NMR and exchangeable protons in ^2H NMR. *Chem. Commun.* **2004**, 1330-1331.
1582. Leyes AE, Poulter CD. Synthesis of (R)-[2- ^2H]isopentenyl diphosphate and determination of its enantiopurity by ^2H NMR spectroscopy in a lyotropic medium. *Org. Lett.* **1999**, 1, 1067-1070.
1583. Razon P, Dhulut S, Bezzenine-Lafollée S, Courtieu J, Pancrazi A, Ardisson J. Secondary allyltitanium(IV) reagents in aldehyde allylation I. Extension of the Hoppe reaction to γ -alkoxy secondary allyl carbamates. *Synthesis* **2005**, 102-108.
1584. Goss RJM, Meddour A, Courtieu J, O'Hagan D. [$^2\text{H}_1$]-Fluoroacetic acid. Resolution and assignment of the absolute configuration by ^2H -NMR in chiral liquid crystalline solvent. *Synthesis and Applications of Isotopically Labelled Compounds* **2001**, 7, 163-166.
1585. Lesot P, Sarfati M, Merlet D, Ancian B, Emsley JW, Timimi BA. 2D-NMR strategy dedicated to the analysis of weakly ordered, fully deuterated enantiomers in chiral liquid crystals. *J. Am. Chem. Soc.* **2003**, 125, 7689-7695.
1586. Lafon O, Berdagué P, Lesot P. Use of two-dimensional correlation between ^2H quadrupolar splittings and ^{13}C CSA's for assignment of NMR spectra in chiral nematics. *Phys. Chem. Chem. Phys.* **2004**, 6, 1080-1084.
1587. Merlet D, Ancian B, Smadja W, Courtieu J, Lesot P. Analysis of natural abundance deuterium NMR spectra of enantiomers in chiral liquid crystals via 2D auto-correlation experiments. *Chem. Commun.* **1998**, 2301-2302.
1588. Lesot P, Merlet D, Lowenstein A, Courtieu J. Enantiomeric visualization using proton-decoupled natural abundance deuterium NMR in poly (γ -benzyl-L-glutamate) liquid crystalline solutions. *Tetrahedron: Asymmetry* **1998**, 9, 1871-1881.
1589. Merlet D, Ancian B, Courtieu J, Lesot P. Two-dimensional deuterium NMR spectroscopy of chiral molecules oriented in a polypeptide liquid crystal: applications for the enantiomeric analysis through natural abundance deuterium NMR. *J. Am. Chem. Soc.* **1999**, 121, 5249-5258.
1590. Sarfati M, Courtieu J, Lesot P. First successful enantiomeric discrimination of chiral alkanes using NMR spectroscopy. *Chem. Commun.* **2002**, 1113-1114.
1592. Lafon O, Lesot P, Merlet D, Courtieu J. Modified z-gradient filtering as a mean to obtain phased deuterium autocorrelation 2D NMR spectra in oriented solvents. *J. Magn. Reson.* **2004**, 171, 135-142.
1593. Parenty A, Campagne J, Aroulanda C, Lesot P. Routine use of natural abundance deuterium NMR in a polypeptidic chiral oriented solvent for the determination of the enantiomeric composition of chiral building blocks. *Org. Lett.* **2002**, 4, 1663-1666.
1594. Lesot P, Sarfati M, Courtieu J. Natural abundance deuterium NMR spectroscopy in polypeptide liquid crystals as a new and incisive means for the enantiodifferentiation of chiral hydrocarbons. *Chem. Eur. J.* **2003**, 9, 1724-1745.
1595. Aroulanda C, Lesot P, Merlet D, Courtieu J. Structural ambiguities revisited in two bridged ring systems exhibiting enantiotopic elements, using natural abundance deuterium NMR in chiral liquid crystals. *J. Phys. Chem. A.* **2003**, 107, 10911-10918.
1596. Lesot P, Aroulanda C, Billault I. Exploring the analytical potential of NMR spectroscopy in chiral anisotropic media for the study of the natural abundance deuterium distribution in organic molecules. *Anal. Chem.* **2004**, 76, 2827-2835.